

International Journal of Modern Physics A
 © World Scientific Publishing Company

NEUTRINO MIXING WITH NON-ZERO θ_{13} IN ZEE-BABU MODEL

HOANG NGOC LONG

Institute of Physics, VAST, 10 Dao Tan, Ba Dinh, Hanoi, Vietnam
hnlng@iop.vast.ac.vn

VO VAN VIEN

Department of Physics, Tay Nguyen University, 567 Le Duan, Buon Ma Thuot, Vietnam
vvienk16@gmail.com

Received Day Month Year
 Revised Day Month Year

The exact solution for the neutrino mass matrix of the Zee-Babu model is derived. Tribi-maximal mixing imposes conditions on the Yukawa couplings, from which the normal mass hierarchy is preferred. The derived conditions give a possibility of Majorana maximal CP violation in the neutrino sector. We have shown that non-zero θ_{13} is generated if Yukawa couplings between leptons almost equal to each other. The model gives some regions of the parameters where neutrino mixing angles and the normal neutrino mass hierarchy obtained consistent with the recent experimental data.

Keywords: Neutrino mass and mixing, Non-standard-model neutrinos, Zee-Babu model

PACS numbers: 14.60.Pq, 14.60.St

1. Introduction

Nowadays, particle physicists are attracted by two exciting subjects: Higgs and neutrino physics. The neutrino mass and mixing are the first evidence of beyond Standard Model physics. Many experiments show that neutrinos have tiny masses and their mixing is still mysterious.^{1,2} Recent data are a clear sign of rather large value θ_{13} .³

The tribimaximal (TBM) form for explaining the lepton mixing scheme was first proposed by Harrison-Perkins-Scott (HPS), which apart from the phase redefinitions, is given by^{4,5}

$$U_{\text{HPS}} = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad (1)$$

can be considered as a good approximation for the recent neutrino experimental data, where the large mixing angles are completely different from the quark mixing ones defined by the Cabibbo-Kobayashi-Maskawa (CKM) matrix⁶⁻⁸.

The most recent fits suggest that one of the mixing angles is approximately zero and another has a value that implies a mass eigenstate that is nearly an equal mixture of ν_μ and ν_τ . The parameters of neutrino oscillations such as the squared mass differences and mixing angles are now very constrained. The data in PDG2010⁹ imply

$$\sin^2(2\theta_{12}) = 0.87 \pm 0.03, \quad \sin^2(2\theta_{23}) > 0.92, \quad \sin^2(2\theta_{13}) < 0.15, \\ \Delta m_{21}^2 = (7.59 \pm 0.20) \times 10^{-5} \text{eV}^2, \quad \Delta m_{32}^2 = (2.43 \pm 0.13) \times 10^{-3} \text{eV}^2, \quad (2)$$

where (and hereafter) the best fits are taken into accounts. Whereas, the new data^{10–19} have been given to be slightly modified from the old fits (2):

$$\sin^2(2\theta_{12}) = 0.857 \pm 0.024, \quad \sin^2(2\theta_{13}) = 0.098 \pm 0.013, \quad \sin^2(2\theta_{23}) > 0.95, \\ \Delta m_{21}^2 = (7.50 \pm 0.20) \times 10^{-5} \text{eV}^2, \quad \Delta m_{32}^2 = (2.32^{+0.12}_{-0.08}) \times 10^{-3} \text{eV}^2. \quad (3)$$

On the other hand, the discovery of the the long- awaited Higgs boson at around 125 GeV²⁰ opened a new chapter in particle physics. It is essential for us to determine which model the discovered Higgs boson belongs to? For this aim, the diphoton decay of the Higgs boson plays a very important role. It is expected that new physics might enter here to modify the standard model (SM) Higgs property.

For the above mentioned reasons, the search for an extended model coinciding with the current data on neutrino and Higgs physics is one of top our priorities. In our opinion, the model with the simplest particle content is preferred. In the SM, neutrinos are strictly massless. For neutrino mass, an original model pointed out by Zee in Ref.²¹ in which new scalars are added in the Higgs sector with neutrino masses induced at the one-loop level. After that a two-loop scenario called the Zee-Babu model²² was proposed. The Zee-Babu model^{21–23} with just two additional charged Higgs bosons (h^-, k^{--}) carrying lepton number 2, is very attractive^a. In this model, neutrinos get mass from two-loop radiative corrections, which can fit current neutrino data. Moreover, the singly and doubly charged scalars that are new in the model can explain the large annihilation cross section of a dark matter pair into two photons as hinted-at by the recent analysis of the Fermi γ -ray space telescope data²⁴, if the new charged scalars are relatively light and have large couplings to a pair of dark matter particles. These new scalars can also enhance the $B(H \rightarrow \gamma\gamma)$, as the recent LHC results may suggest.

The Zee-Babu model contains the Yukawa couplings which are specific for lepton number violating processes. There has been much work^{26–30} constraining the parameter space of the model, however the explicit values of neutrino masses and mixings have not been considered.

In this paper, starting from the neutrino mass matrix, we get the exact solution, i.e., the eigenstates and the eigenvalues. As a consequence, the neutrino mixing matrix follows. With this exact solution, we can fit current data and get constraints

^aIn the recent paper,²⁹ the parameter space of the model under consideration has been reanalyzed, and the lower bounds for masses of the singly and doubly charged Higgses lie between 1 to 2 TeV.




Fig. 1. The two-loop diagram in the Zee-Babu model.

on the couplings. We hope that experiments in the near future will approve or rule out the model.

2. Neutrino mass matrix in the Zee-Babu model

The Zee-Babu model²² includes two $SU(2)_L$ singlet Higgs fields, a singly charged field h^- and a doubly charged field k^{--} . Moreover, right-handed neutrinos are not introduced. The addition of these singlets gives rise to the Yukawa couplings:

$$\mathcal{L}_Y = f_{ab} \overline{(\psi_{aL})^C} \psi_{bL} h^+ + h'_{ab} \overline{(l_{aR})^C} l_{bR} k^{++} + H.c., \quad (4)$$

where ψ_L stands for the left-handed lepton doublet, l_R for the right-handed charged lepton singlet and $(a, b = e, \mu, \tau)$ being the generation indices, a superscript C indicating charge conjugation. Here $\psi^C = C \overline{\psi}^T$ with C being the charge-conjugation matrix. The coupling constant f_{ab} is antisymmetric ($f_{ab} = -f_{ba}$), whereas h_{ab} is symmetric ($h_{ab} = h_{ba}$). Gauge invariance precludes the singlet Higgs fields from coupling to the quarks. In terms of the component fields, the interaction Lagrangian is given by

$$\begin{aligned} \mathcal{L}_Y = 2 [& f_{e\mu} (\bar{\nu}_e^c \mu_L - \bar{\nu}_\mu^c e_L) + f_{e\tau} (\bar{\nu}_e^c \tau_L - \bar{\nu}_\tau^c e_L) + f_{\mu\tau} (\bar{\nu}_\mu^c \tau_L - \bar{\nu}_\tau^c \mu_L)] h^+ \\ & + [h_{ee} \bar{e}^c e_R + h_{\mu\mu} \bar{\mu}^c \mu_R + h_{\tau\tau} \bar{\tau}^c \tau_R + h_{e\mu} \bar{e}^c \mu_R + h_{e\tau} \bar{e}^c \tau_R + h_{\mu\tau} \bar{\mu}^c \tau_R] k^{++} \quad (5) \\ & + H.c. \end{aligned}$$

where we have used $h_{aa} = h'_{aa}$, $h_{ab} = 2h'_{ab}$ for $a \neq b$. Eq. (4) conserves lepton number, therefore, itself cannot be responsible for neutrino mass generation.

The Higgs potential contains the terms:

$$V(\phi, h^+, k^{++}) = \mu(h^- h^- k^{++} + h^+ h^+ k^{--}) + \dots, \quad (6)$$

which violate lepton number by two units. They are expected to cause Majorana neutrino masses.

In the literature, Majorana neutrino masses are generated at the two-loop level via the diagram shown in²³ and again depicted in Fig.1. The corresponding mass

matrix for Majorana neutrinos is as follows

$$M_{ab} = 8\mu f_{ac} h_{cd}^* m_c m_d I_{cd}(f^+)_{db}. \quad (7)$$

The integral I_{cd} is given by³¹

$$I_{cd} = \int \frac{d^4 k}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} \frac{1}{k^2 - m_c^2} \frac{1}{k^2 - M_h^2} \frac{1}{q^2 - m_d^2} \times \frac{1}{q^2 - M_h^2} \frac{1}{(k - q)^2 - M_k^2} \quad (8)$$

Note that Eq. (8) can be simplified by neglecting the charged lepton masses m_c and m_d ²⁵.

To evaluate the integral above, one neglects the charged lepton masses in the denominator, since these masses are much smaller than the charged scalar masses M_h and M_k . Then

$$I_{cd} \simeq I = \frac{1}{(16\pi^2)^2} \frac{1}{M^2} \frac{\pi^2}{3} \tilde{I}(r), \quad M \equiv \max(M_k, M_h) \quad (9)$$

which does not depend on lepton masses. Here $\tilde{I}(r)$ is a function of the ratio of the masses of the charged Higgses $r \equiv M_k^2/M_h^2$,

$$\tilde{I}(r) = \begin{cases} 1 + \frac{3}{\pi^2}(\log^2 r - 1) & \text{for } r \gg 1 \\ 1 & \text{for } r \rightarrow 0, \end{cases} \quad (10)$$

which is close to 1 for a wide range of scalar masses.

The neutrino mass matrix arising from (7) is symmetric and given by

$$\mathcal{M}_\nu = -I\mu f_{\mu\tau}^2 \times \begin{pmatrix} \epsilon^2\omega_{\tau\tau} + 2\epsilon\epsilon'\omega_{\mu\tau} + \epsilon'^2\omega_{\mu\mu} & \epsilon\omega_{\tau\tau} + \epsilon'(\omega_{\mu\tau} - \epsilon\omega_{e\tau} - \epsilon'\omega_{e\mu}) & -\epsilon'\omega_{\mu\mu} - \epsilon(\omega_{\mu\tau} + \epsilon\omega_{e\tau} + \epsilon'\omega_{e\mu}) \\ \star & \omega_{\tau\tau} + \epsilon'^2\omega_{ee} - 2\epsilon'\omega_{e\tau} & \epsilon\epsilon'\omega_{ee} - \omega_{\mu\tau} - \epsilon\omega_{e\tau} + \epsilon'\omega_{e\mu} \\ \star & \star & \omega_{\mu\mu} + 2\epsilon\omega_{e\mu} + \epsilon'^2\omega_{ee} \end{pmatrix} \quad (11)$$

where we have redefined parameters:

$$\epsilon \equiv \frac{f_{e\tau}}{f_{\mu\tau}}, \quad \epsilon' \equiv \frac{f_{e\mu}}{f_{\mu\tau}}, \quad \omega_{ab} \equiv m_a h_{ab}^* m_b. \quad (12)$$

Let us denote

$$\begin{aligned} \omega'_{\tau\tau} &\equiv \omega_{\tau\tau} + \epsilon'^2\omega_{ee} - 2\epsilon'\omega_{e\tau}, \\ \omega'_{\mu\tau} &\equiv \omega_{\mu\tau} + \epsilon\omega_{e\tau} - \epsilon'\omega_{e\mu} - \epsilon\epsilon'\omega_{ee}, \\ \omega'_{\mu\mu} &\equiv \omega_{\mu\mu} + 2\epsilon\omega_{e\mu} + \epsilon'^2\omega_{ee}, \end{aligned} \quad (13)$$

then the neutrino mass matrix can be rewritten in the compact form

$$\mathcal{M}_\nu = -I\mu f_{\mu\tau}^2 \begin{pmatrix} \epsilon^2\omega'_{\tau\tau} + 2\epsilon\epsilon'\omega'_{\mu\tau} + \epsilon'^2\omega'_{\mu\mu} & \epsilon\omega'_{\tau\tau} + \epsilon'\omega'_{\mu\tau} - \epsilon\omega'_{\mu\tau} - \epsilon'\omega'_{\mu\mu} \\ \star & \omega'_{\tau\tau} & -\omega'_{\mu\tau} \\ \star & \star & \omega'_{\mu\mu} \end{pmatrix}. \quad (14)$$

The above matrix has three exact eigenvalues given by

$$m_1 = 0, \\ m_{2,3} = \frac{1}{2} \left(-kF \pm \sqrt{k^2 [F^2 + 4(1 + \epsilon^2 + \epsilon'^2)(\omega'_{\mu\tau}{}^2 - \omega'_{\mu\mu}\omega'_{\tau\tau})]} \right), \quad (15)$$

where we have denoted

$$k = \mu I f_{\mu\tau}^2, \quad F = (1 + \epsilon'^2)\omega'_{\mu\mu} + 2\epsilon\epsilon'\omega'_{\mu\tau} + (1 + \epsilon^2)\omega'_{\tau\tau}. \quad (16)$$

The massless eigenstate is given by

$$\nu_1 = \frac{1}{\sqrt{f_{e\mu}^2 + f_{e\tau}^2 + f_{\mu\tau}^2}} (f_{\mu\tau}\nu_e - f_{e\tau}\nu_\mu + f_{e\mu}\nu_\tau). \quad (17)$$

The mass matrix (14) is diagonalized as

$$U^T \mathcal{M}_\nu U = \text{diag}(0, m_2, m_3),$$

where

$$U = \begin{pmatrix} \frac{1}{\sqrt{1+\epsilon^2+\epsilon'^2}} & -\frac{A_1}{\sqrt{1+A_1^2+B_1^2}} & \frac{A_2}{\sqrt{1+A_2^2+B_2^2}} \\ -\frac{\epsilon}{\sqrt{1+\epsilon^2+\epsilon'^2}} & -\frac{B_1}{\sqrt{1+A_1^2+B_1^2}} & \frac{B_2}{\sqrt{1+A_2^2+B_2^2}} \\ \frac{\epsilon'}{\sqrt{1+\epsilon^2+\epsilon'^2}} & -\frac{1}{\sqrt{1+A_1^2+B_1^2}} & \frac{1}{\sqrt{1+A_2^2+B_2^2}} \end{pmatrix} \quad (18)$$

with the new notations

$$A_{1,2} = \frac{-k [\epsilon(\epsilon'^2 - 1)\omega'_{\mu\mu} + 2\epsilon'(1 + \epsilon^2)\omega'_{\mu\tau} + \epsilon(1 + \epsilon^2)\omega'_{\tau\tau}] \pm \epsilon\sqrt{k^2 F'}}{2k [\epsilon\epsilon'\omega'_{\mu\mu} + (1 + \epsilon^2)\omega'_{\mu\tau}]}, \quad (19)$$

$$B_{1,2} \equiv \frac{k(1 + \epsilon'^2)\omega'_{\mu\mu} - k(1 + \epsilon^2)\omega'_{\tau\tau} \pm \sqrt{k^2 F'}}{2k [\epsilon\epsilon'\omega'_{\mu\mu} + (1 + \epsilon^2)\omega'_{\mu\tau}]}, \quad (20)$$

and

$$F' = F^2 + 4(1 + \epsilon^2 + \epsilon'^2)(\omega'_{\mu\tau}{}^2 - \omega'_{\mu\mu}\omega'_{\tau\tau}). \quad (21)$$

The eigenstates ν_i corresponding to the eigenvalues m_i ($i = 1, 2, 3$) are found to be

$$\nu_1 = \frac{1}{\sqrt{f_{e\mu}^2 + f_{e\tau}^2 + f_{\mu\tau}^2}} (f_{\mu\tau}\nu_e - f_{e\tau}\nu_\mu + f_{e\mu}\nu_\tau), \\ \nu_2 = -\frac{A_1}{\sqrt{1+A_1^2+B_1^2}}\nu_e - \frac{B_1}{\sqrt{1+A_1^2+B_1^2}}\nu_\mu - \frac{1}{\sqrt{1+A_1^2+B_1^2}}\nu_\tau, \\ \nu_3 = \frac{A_2}{\sqrt{1+A_2^2+B_2^2}}\nu_e + \frac{B_2}{\sqrt{1+A_2^2+B_2^2}}\nu_\mu + \frac{1}{\sqrt{1+A_2^2+B_2^2}}\nu_\tau. \quad (22)$$

From the explicit expressions of $A_{1,2}$ and $B_{1,2}$ in (19) and (20), some useful relations are in order

$$\begin{aligned} A_1 A_2 + B_1 B_2 + 1 &= 0, \\ A_1 - \epsilon B_1 + \epsilon' &= 0, \\ A_2 - \epsilon B_2 + \epsilon' &= 0, \\ (A_1 - A_2)/(B_1 - B_2) &= \epsilon. \end{aligned} \quad (23)$$

One also has

$$\begin{aligned} A_1 A_2 &= \frac{(\epsilon'^2 - \epsilon^2)\omega'_{\mu\tau} + \epsilon\epsilon'(\omega'_{\tau\tau} - \omega'_{\mu\mu})}{\epsilon\epsilon'\omega'_{\mu\mu} + (1 + \epsilon^2)\omega'_{\mu\tau}}, \\ B_1 B_2 &= -\frac{(1 + \epsilon'^2)\omega'_{\mu\tau} + \epsilon\epsilon'\omega'_{\tau\tau}}{\epsilon\epsilon'\omega'_{\mu\mu} + (1 + \epsilon^2)\omega'_{\mu\tau}}. \end{aligned}$$

3. Constraints from the tribimaximal mixing

The current data on neutrino mass and mixing show that tribimaximal mixing^{4,5} as displayed in (1) is very specific. Comparing (18) with (1) yields the following conditions

$$\epsilon = \epsilon' = \frac{1}{2}, \quad (24)$$

$$A_2 = 0, \quad A_1 = B_1 = -1, \quad B_2 = 1 \quad (25)$$

Eqs. (24) and (12) lead to

$$f_{e\mu} = f_{e\tau} = \frac{1}{2}f_{\mu\tau} \quad (26)$$

Substitution of (24) into expressions of $A_{1,2}$, $B_{1,2}$ in (19) and (20) yields

$$A_{1,2} = \frac{k(3\omega'_{\mu\mu} - 10\omega'_{\mu\tau} - 5\omega'_{\tau\tau}) \pm \sqrt{k^2 F_0}}{4k(\omega'_{\mu\mu} + 5\omega'_{\mu\tau})}, \quad (27)$$

$$B_{1,2} = \frac{5k(\omega'_{\mu\mu} - \omega'_{\tau\tau}) \pm \sqrt{k^2 F_0}}{2k(\omega'_{\mu\mu} + 5\omega'_{\mu\tau})}, \quad (28)$$

with

$$F_0 = 4(\omega'_{\mu\mu} + 5\omega'_{\mu\tau})^2 + (\omega'_{\mu\mu} - \omega'_{\mu\tau})(21\omega'_{\mu\mu} - 20\omega'_{\mu\tau} - 25\omega'_{\tau\tau}). \quad (29)$$

If $\omega'_{\mu\mu} = \omega'_{\tau\tau} = \omega'$ we have:

$$A_{1,2} = -\frac{1}{2} \left(1 \mp \frac{k(\omega' + 5\omega'_{\mu\tau})}{\sqrt{k^2(\omega' + 5\omega'_{\mu\tau})^2}} \right), \quad (30)$$

$$B_{1,2} = \pm \frac{k(\omega' + 5\omega'_{\mu\tau})}{\sqrt{k^2(\omega' + 5\omega'_{\mu\tau})^2}}. \quad (31)$$

It can be checked that with the help of (24), all remaining conditions in (25) are satisfied if

$$\omega'_{\mu\mu} = \omega'_{\tau\tau} \equiv \omega', \quad (32)$$

and $k(\omega' + 5\omega'_{\mu\tau})$ are negative real numbers. This can be equivalently converted into a relation among the Yukawa couplings

$$\omega_{\mu\mu} + \omega_{e\mu} = \omega_{\tau\tau} - \omega_{e\tau} \quad (33)$$

Note that our derived constraints are somewhat different from those given in²⁷.

From the conditions (24) and (32) we obtain^b

$$m_1 = 0, \quad m_{2,3} = -\frac{1}{4} \left[k(5\omega' + \omega'_{\mu\tau}) \mp \sqrt{k^2(\omega' + 5\omega'_{\mu\tau})^2} \right]. \quad (34)$$

The complex phases which can arise when diagonalizing the neutrino mass matrix (14) can be absorbed by the redefinition of the mass matrix eigenvectors, as it should be given that both $m_{2,3}$ are physical observables. Hence, in this work we assume m_2, m_3 to be real.

Depending on the sign of the function in the square root we have two cases in which $k(5\omega' + \omega'_{\mu\tau})$ being either positive or negative. To fit the experimental data in⁹ the following condition must be satisfied

$$k(\omega' + 5\omega'_{\mu\tau}) < 0. \quad (35)$$

The neutrino masses in (15) becomes

$$m_1 = 0, \quad m_2 = -\frac{3k}{2}(\omega' + \omega'_{\mu\tau}), \quad m_3 = k(-\omega' + \omega'_{\mu\tau}). \quad (36)$$

Taking the central values from the data⁹ as displayed in (2), we have the two following solutions:

(1) $m_1 = 0, \quad m_2 = 0.008712 \text{ eV}, m_3 = -0.050059 \text{ eV}$ and then

$$U = \begin{pmatrix} \frac{2}{\sqrt{6}} & 0.57735 & 4.17428 \times 10^{-17} \\ -\frac{1}{\sqrt{6}} & 0.57735 & 0.707107 \\ \frac{1}{\sqrt{6}} & -0.57735 & 0.707107 \end{pmatrix}. \quad (37)$$

In this case, $\omega'_{\mu\tau}$ and ω' depend only on k due to the following relations:

$$\omega'_{\mu\tau} = -\frac{0.0279335}{k}, \quad \omega' = \frac{0.0221255}{k}, \quad (38)$$

$$\frac{\omega'_{\mu\tau}}{\omega'} = -1.2625, \quad (39)$$

$$k(\omega' + 5\omega'_{\mu\tau}) = -0.117542 < 0.$$

^bThe integration in Fig.1 is linear divergent and has a surface term³², which give a similar form of mass matrix.

(2) $m_1 = 0$, $m_2 = -0.00871206 \text{ eV}$, $m_3 = -0.050059 \text{ eV}$, and

$$U = \begin{pmatrix} \frac{2}{\sqrt{6}} & 0.57735 & -5.93338 \times 10^{-17} \\ -\frac{1}{\sqrt{6}} & 0.57735 & -0.707107 \\ -\frac{1}{\sqrt{6}} & 0.57735 & 0.707107 \end{pmatrix}. \quad (40)$$

In this case $\omega'_{\mu\tau}$ and ω' depend only on k according to the following relations:

$$\omega'_{\mu\tau} = -\frac{0.0221255}{k}, \quad \omega' = \frac{0.0279335}{k}, \quad (41)$$

$$\frac{\omega'_{\mu\tau}}{\omega'} = -0.792076, \quad (42)$$

$$k(\omega' - 5\omega'_{\mu\tau}) = -0.0826938 < 0.$$

The expressions (39) and (42) show that $\omega'_{\mu\mu}$, $\omega'_{\tau\tau}$ and $\omega'_{\mu\tau}$ are of the same order, and the normal neutrino mass hierarchy was used^c.

In terms of the usual neutrino-oscillation parameters, the matrices (37) and (40) mean that

$$\sin^2_{23} = \frac{1}{2}, \quad \sin^2_{12} = \frac{1}{3}, \quad \sin^2_{13} = 0. \quad (43)$$

which in good agreement with the tribimaximal form.⁹ However, with a vanishing θ_{13} now excluded at more than $10\sigma^{17}$ the situation has changed somewhat and the result in (2) should be considered just as a good approximation.

Using the standard parametrization of the neutrino mixing matrix (the PMNS matrix) in terms of three angles and CP violating phases⁶⁻⁸

$$\begin{aligned} U &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\gamma/2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\gamma/2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned} \quad (44)$$

where δ and γ are the Dirac and Majorana CP phase, respectively, and $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$ ($ij = 12, 23, 13$). The above Majorana mass matrix is diagonalized by the PMNS matrix

$$U^T \mathcal{M}_\nu U = \mathcal{M}_{diag} = \text{diag}(m_1, m_2, m_3).$$

^cHere, we have assumed a normal neutrino mass hierarchy in which $m_1 = \lambda_1 = 0, m_2 = \lambda_2, m_3 = \lambda_3$ where λ_i ($i = 1, 2, 3$) are eigenvalues of M_ν in (14). A spectrum with inverted ordering can be obtained by using the notation $m'_3 = \lambda_1 = 0, m'_2 = \lambda_3 \equiv m_3$ and $m'_1 = \lambda_2 \equiv m_2$.

m_2 [eV]	m_3 [eV]	$e^{-i\gamma}$	γ [rad]
0.008712	-0.0500591	1.00002	$\pi/2$
0.00871	-0.0500591	1.00001	$\pi/2$

In the case of the normal mass hierarchy, the four parameters are described as^{25,27}

$$\begin{aligned}\epsilon &= \tan \theta_{12} \frac{s_{23}}{c_{13}} + \tan \theta_{13} e^{i\delta}, \\ \epsilon' &= \tan \theta_{12} \frac{s_{23}}{c_{13}} - \tan \theta_{13} e^{i\delta}, \\ \frac{\omega'_{\mu\tau}}{\omega'_{\mu\mu}} &= -\frac{c_{13}^2 s_{23} c_{23}}{c_{13}^2 c_{23}^2 + r_{2/3}(s_{12} s_{13} c_{23} e^{-i\delta} + c_{12} s_{23})^2 e^{-i\gamma}} \\ &\quad - \frac{r_{2/3}(s_{12} s_{13} c_{23} e^{-i\delta} + c_{12} s_{23})(s_{12} s_{13} s_{23} e^{-i\delta} - c_{12} c_{23}) e^{-i\gamma}}{c_{13}^2 c_{23}^2 + r_{2/3}(s_{12} s_{13} c_{23} e^{-i\delta} + c_{12} s_{23})^2 e^{-i\gamma}},\end{aligned}\quad (45)$$

$$\frac{\omega'_{\tau\tau}}{\omega'_{\mu\mu}} = \frac{c_{13}^2 s_{23}^2 + r_{2/3}(s_{12} s_{13} s_{23} e^{-i\delta} - c_{12} c_{23})^2 e^{-i\gamma}}{c_{13}^2 c_{23}^2 + r_{2/3}(s_{12} s_{13} c_{23} e^{-i\delta} + c_{12} s_{23})^2 e^{-i\gamma}}, \quad (46)$$

with $r_{2/3} = m_2/m_3$.

We can easily see that with the help of (32), Eq. (46) is automatically satisfied. On the other hand, from (45) ones can find the values of γ corresponding to those of m_2, m_3 as shown in Table 1, in which the values of γ is approximately equal to $\frac{\pi}{2}$. So the condition (32) leads to Majorana maximal CP violation: $\sin \gamma_{CP} \simeq 1$, as mentioned in Ref. 34.

The recent considerations have implied $\theta_{13} \neq 0$, but small as given in Ref.10. A deviations from the tribimaximal form would be achieved with a non-zero value of A_2 and a small difference of ϵ and ϵ' as shown in Section 4.

4. Experimental constraints with non-zero θ_{13}

The realistic neutrino mixing will be slightly deviated from the tribimaximal form. This will be achieved with a very small value of A_2 and $\epsilon' \simeq \epsilon \simeq \frac{1}{2}$. With the help of (23), the matrix U in (18) becomes

$$U = \begin{pmatrix} \frac{1}{\sqrt{1+\epsilon^2+\epsilon'^2}} & \frac{\epsilon^2+\epsilon'(\epsilon'+A_2)}{\sqrt{(1+\epsilon^2+\epsilon'^2)[\epsilon^2+(1+\epsilon^2)A_2^2+2A_2\epsilon'+\epsilon'^2]}} & \frac{A_2\epsilon}{\sqrt{(1+A_2^2)\epsilon^2+(A_2+\epsilon')^2}} \\ -\frac{\epsilon}{\sqrt{1+\epsilon^2+\epsilon'^2}} & \frac{\epsilon(1-A_2\epsilon')}{\sqrt{(1+\epsilon^2+\epsilon'^2)[\epsilon^2+(1+\epsilon^2)A_2^2+2A_2\epsilon'+\epsilon'^2]}} & \frac{A_2+\epsilon'}{\sqrt{(1+A_2^2)\epsilon^2+(A_2+\epsilon')^2}} \\ \frac{\epsilon'}{\sqrt{1+\epsilon^2+\epsilon'^2}} & -\frac{A_2(1+\epsilon^2)+\epsilon'}{\sqrt{(1+\epsilon^2+\epsilon'^2)[\epsilon^2+(1+\epsilon^2)A_2^2+2A_2\epsilon'+\epsilon'^2]}} & \frac{\epsilon}{\sqrt{(1+A_2^2)\epsilon^2+(A_2+\epsilon')^2}} \end{pmatrix} \quad (47)$$

Combining (47) and (44) we obtain:

$$t_{12} = \frac{U_{12}}{U_{11}} = \frac{\epsilon^2 + \epsilon'(A_2 + \epsilon')}{\sqrt{\epsilon^2 + (1 + \epsilon^2)A_2^2 + 2A_2\epsilon' + \epsilon'^2}}, \quad (48)$$

$$t_{23} = \frac{U_{23}}{U_{33}} = \frac{A_2 + \epsilon'}{\epsilon}. \quad (49)$$

with $t_{ij} = \tan \theta_{ij}$ ($ij = 12, 23, 13$).

Since ϵ and ϵ' close to each other, it can be assumed that

$$\epsilon' = \alpha\epsilon \quad (50)$$

where α is a constant close to 1.

From the expressions (48), (49) and (50) we obtain the following relations:

$$t_{23} = -\frac{\alpha\epsilon^3(1+t_{12}^2) + \sqrt{\Gamma}}{\alpha^2\epsilon^3 - \epsilon(1+\epsilon^2)t_{12}^2}, \quad (51)$$

$$A_2 = \frac{\epsilon^3\alpha(1+\alpha^2) - \alpha\epsilon t_{12}^2 + \sqrt{\Gamma}}{t_{12}^2(1+\epsilon^2) - \alpha^2\epsilon^2}, \quad (52)$$

or

$$t_{23} = \frac{-\alpha\epsilon^3(1+t_{12}^2) + \sqrt{\Gamma}}{\alpha^2\epsilon^3 - \epsilon(1+\epsilon^2)t_{12}^2}. \quad (53)$$

$$A_2 = \frac{\epsilon^3\alpha(1+\alpha^2) - \alpha\epsilon t_{12}^2 - \sqrt{\Gamma}}{t_{12}^2(1+\epsilon^2) - \alpha^2\epsilon^2}, \quad (54)$$

where

$$\Gamma = \epsilon^2 t_{12}^2 [1 + (1 + \alpha^2)\epsilon^2][(1 + \alpha^2)\epsilon^2 - t_{12}^2]. \quad (55)$$

Substituting A_2 from (52) into (47) yields

$$U = \begin{pmatrix} U_{11} & U_{12} & U_{13} \\ U_{21} & U_{22} & U_{23} \\ U_{31} & U_{32} & U_{33} \end{pmatrix}, \quad (56)$$

with

$$\begin{aligned} U_{11} &= \frac{1}{\sqrt{1 + (1 + \alpha^2)\epsilon^2}}, & U_{21} &= -\frac{\epsilon}{\sqrt{1 + (1 + \alpha^2)\epsilon^2}} \\ U_{31} &= \frac{\alpha\epsilon}{\sqrt{1 + (1 + \alpha^2)\epsilon^2}}, & U_{12} &= -\frac{\epsilon [\epsilon t_{12}^2 + (1 + \alpha^2)\epsilon^3 t_{12}^2 + \alpha\sqrt{\Gamma}]}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, \\ U_{22} &= \frac{\epsilon [\alpha^4 \epsilon^4 - (1 + \epsilon^2)t_{12}^2 + \alpha^2 \epsilon^2 (1 + \epsilon^2 - t_{12}^2) + \alpha\epsilon\sqrt{\Gamma}]}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, \\ U_{32} &= \frac{\alpha(1 + \alpha^2)\epsilon^5 + \alpha\epsilon^3 + (1 + \epsilon^2)\sqrt{\Gamma}}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, & U_{13} &= \frac{\epsilon [\alpha(1 + \alpha^2)\epsilon^3 - \alpha\epsilon t_{12}^2 + \sqrt{\Gamma}]}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, \\ U_{23} &= -\frac{\alpha\epsilon^3(1 + t_{12}^2) + \sqrt{\Gamma}}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, & U_{33} &= \frac{\epsilon [\alpha^2 \epsilon^2 - t_{12}^2(1 + \epsilon^2)]}{\sqrt{\epsilon^3 [1 + (1 + \alpha^2)\epsilon^2]^2 \Gamma'}}, \end{aligned} \quad (57)$$

where

$$\Gamma' = (1 - \alpha^2)\epsilon t_{12}^2 + \epsilon^3(1 + \alpha^2)(\alpha^2 + t_{12}^2) + 2\alpha\sqrt{\Gamma}. \quad (58)$$

We see that the neutrino mixing matrix in (56) with the elements given in (57) depends only on three parameters α, ϵ and t_{12} . It is easily to show that the model

can fit the recent experimental constraints on the neutrino mixing angles. Indeed, by choosing $\alpha \in (0.98, 1.00)$, $\epsilon \in (0.50, 0.505)$ and taking the new data given in¹⁰ with $t_{12} = 0.691$, we obtain

$$\begin{aligned} U_{11} &\in (0.814 \div 0.818), & U_{12} &\in -(0.563 \div 0.566), & U_{13} &\in (0.010 \div 0.140), \\ U_{21} &\in -(0.409 \div 0.412), & U_{22} &\in -(0.380 \div 0.460), & U_{23} &\in -(0.790 \div 0.830), \\ U_{31} &\in (0.4025 \div 0.410), & U_{32} &\in (0.680 \div 0.720), & U_{33} &\in -(0.540 \div 0.600). \end{aligned} \quad (59)$$

or

$$U = \begin{pmatrix} 0.814 \div 0.818 & -(0.563 \div 0.566) & 0.010 \div 0.140 \\ -(0.409 \div 0.412) & -(0.380 \div 0.460) & -(0.790 \div 0.830) \\ 0.4025 \div 0.410 & 0.680 \div 0.720 & -(0.540 \div 0.600) \end{pmatrix}. \quad (60)$$

It is interesting to note that the model-independent parametrization of non-TBM structures based on deviations from the reactor, solar and atmospheric angles³⁵ and on small perturbations of the tribimaximal mixing eigenvectors³⁶ is similar to our approach here. Our set ϵ, α, t_{12} is equivalent to the set $\epsilon_{12}, \epsilon_{23}, \epsilon_{13}$ in Ref.³⁶

The Figs. 2a, 2b, 2c, Figs.3a, 3b, 3c, Figs.4a, 4b and 4c give the dependence of the elements of U matrix on α and ϵ with $t_{12} = 0.691$.




Fig. 2. U_{11}, U_{21}, U_{31} as functions of α and ϵ with $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$

With $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$, from (51) we obtain $t_{23} \in (1.3, 1.59)$ or $\theta_{23} \in (52.43^\circ, 56.31^\circ)$, and $A_2 \in (0.15, 0.25)$ which are shown in Figs. 5a and 5b, respectively. In this case, $s_{13} \in (0.1, 0.14)$ or $\theta_{13} \in (5.74^\circ, 8.05^\circ)$.

Similarly, substituting A_2 from (54) into (47) yields

$$U = \begin{pmatrix} 0.814 \div 0.818 & 0.563 \div 0.566 & -(0.010 \div 0.140) \\ -(0.409 \div 0.412) & 0.69 \div 0.73 & 0.54 \div 0.58 \\ 0.4025 \div 0.410 & -(0.38 \div 0.44) & 0.8 \div 0.83 \end{pmatrix} \quad (61)$$

provided that $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$. In this case, $t_{23} \in (0.65, 0.75)$ or $\theta_{23} \in (45^\circ, 50.19^\circ)$, $s_{13} \in (0.02, 0.08)$ or $\theta_{13} \in (1.15^\circ, 4.6^\circ)$ and $A_2 \in (0.05, 0.15)$. We note that in these regions of the values of α and ϵ , θ_{13} is smaller than that given




Fig. 3. U_{12}, U_{22}, U_{32} as functions of α and ϵ with $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$




Fig. 4. U_{13}, U_{23}, U_{33} as functions of α and ϵ with $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$




Fig. 5. (a) t_{23} as a function of α and ϵ with $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$; (b) A_2 as a function of α and ϵ with $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.505)$

in¹⁰, but the other regions of these parameters will provide a consistent range of θ_{13} , such as, when $\alpha \in (0.98, 1.00)$ and $\epsilon \in (0.50, 0.51)$ then $|s_{13}| \in (0.1, 0.16)$ or $\theta_{13} \in (5.74^\circ, 9.21^\circ)$. This range of θ_{13} satisfies the recent experimental data in¹⁰.

From (51) and (53) we can have the relations of t_{23} and t_{12}, α, ϵ as shown in the Figs. 6a, 6b and 6c, and 7a, 7b and 7c, respectively, in which the values of θ_{23} obtained encompass the best fit values in¹⁰.




Fig. 6. (a) t_{23} as a function of ϵ with $\epsilon \in (0.50, 0.51)$ and $\alpha = 1, t_{12} = 0.691$; (b) t_{23} as a function of t_{12} with $t_{12} \in (0.690, 0.691)$ and $\alpha = 1, \epsilon = 0.49$; (c) t_{23} as a function of α with $\alpha \in (1.0, 1.02)$, $t_{12} = 0.691, \epsilon = 0.50$ from (51).




Fig. 7. (a) t_{23} as a function of ϵ with $\epsilon \in (0.50, 0.505)$ and $\alpha = 1, t_{12} = 0.691$; (b) t_{23} as a function of t_{12} with $t_{12} \in (0.690, 0.691)$ and $\alpha = 1, \epsilon = 0.50$; (c) t_{23} as a function of α with $\alpha \in (1.00, 1.01)$, $t_{12} = 0.691, \epsilon = 0.50$ from (53).

With the help of (50), F in (16) becomes:

$$F = (1 + \alpha^2 \epsilon^2) \omega'_{\mu\mu} + 2\alpha \epsilon^2 \omega'_{\mu\tau} + (1 + \epsilon^2) \omega'_{\tau\tau}, \quad (62)$$

and the physical neutrino masses from (15) is defined

$$m_1 = 0, \\ m_{2,3} = \frac{1}{2} \left(-kF \pm \sqrt{k^2(F^2 + B)} \right), \quad (63)$$

with

$$B = 4[1 + (1 + \alpha^2)\epsilon^2](\omega'^2_{\mu\tau} - \omega'_{\mu\mu}\omega'_{\tau\tau}). \quad (64)$$

Taking the central values from the data¹⁰ on neutrino mass square differences:

$$\Delta m_{21}^2 = 7.50 \times 10^{-5} \text{ eV}^2, \quad \Delta m_{32}^2 = 2.32 \times 10^{-3} \text{ eV}^2, \quad (65)$$

we obtain

$$k = \frac{0.0402359}{F}. \quad (66)$$

The neutrino masses are explicitly given as

$$m_1 = 0, \quad m_2 = 0.00871206 \text{ eV}, \quad m_3 = -0.048948 \text{ eV}, \quad (67)$$

which are in a normal ordering.

The ratio of m_2 to m_3 is given

$$\frac{|m_2|}{|m_3|} = 0.177986 \quad (68)$$

which is the same order as in Ref. 27.

Without loss of generality, we assume $\omega'_{\mu\mu} = \omega'_{\tau\tau} = \omega'$. From (16), (50) and (66) we obtain the dependence of k on α, ϵ and ω and ω'_{23} :

$$k = \frac{0.0402359}{[(2 + (1 + \alpha^2)\epsilon^2)]\omega' + 2\alpha\epsilon^2\omega'_{\mu\tau}}. \quad (69)$$

In the case $\alpha = 1.00$ and $\epsilon = 0.5$, one has

$$k = -\frac{0.0402359}{2.5\omega' + 0.5\omega'_{\mu\tau}}.$$

The Fig. 8 gives the dependence of k on $\omega', \omega'_{\mu\tau}$ with $\omega'_{\mu,\tau} \in (0.95, 1.0)$ and $\omega' \in (0.80, 0.9)$.




Fig. 8. k as a function of $\alpha, \epsilon, \omega'$ and ω'_{23} with $\omega'_{\mu,\tau} \in (0.95, 1.0)$, $\omega' \in (0.80, 0.9)$ and $\alpha = 1.00$, $\epsilon = 0.5$

5. Summary

In this paper we have derived the exact eigenvalues and eigenstates of the neutrino mass matrix in the Zee-Babu model. Tribimaximal mixing imposes some conditions on the Yukawa couplings. The constraints derived in this work slightly differ from other ones given in the literature, and the normal mass hierarchy is preferred. The derived conditions give a possibility of Majorana maximal CP violation in the neutrino sector. We have shown that non-zero θ_{13} is generated if Yukawa couplings between leptons almost equal to each other. We have analyzed behaviors of the mixing angles as functions of the Yukawa couplings, and the model parameter space has been derived.

Acknowledgments

This research is funded by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.01-2011.63.

References

1. G. Altarelli, *An Overview of Neutrino Mixing*, Nucl.Phys.Proc.Suppl. 241-242 (2013) 77-82, arXiv: 1210.3467 [hep-ph].
2. G. Altarelli, *Neutrino Mixing: Theoretical Overview*, ArXiv: 1304.5047 (2013), and references therein.
3. A. Yu. Smirnov, *Neutrino 2012: Outlook - theory*, Nucl.Phys.Proc.Suppl. 235-236 (2013) 431-440, arXiv: 1210.4061 [hep-ph].
4. P. F. Harrison, D. H. Perkins and W. G. Scott, *Phys. Lett. B* **530**, 167 (2002).
5. P. F. Harrison and W. G. Scott, *Phys. Lett. B* **535**, 163 (2002).
6. Z. Maki, M. Nakagawa and S. Sakata, *Prog. Theor. Phys.* **28**, 870 (1962).
7. B. Pontecorvo, *Zh. Eksp. Teor. Fiz.* **53**, 1717 (1967).
8. B. Pontecorvo, *Sov. Phys. JETP* **26**, 984(1968).
9. K. Nakamura *et al.*, Particle Data Group, *J. Phys. G* **37**, 075021 (2010).
10. J. Beringer, *et al.*(2012), *Review of Particle Physics* (Particle Data Group), Phys. Rev. D. 86, 010001.
11. T. Schwetz, M. Tortola, and J. Valle (2011), New J. Phys. 13, 109401, arXiv: 131108.1376 [hep-ph].
12. K. Abe *et al.*(2011) [T2K Collaboration], Phys. Rev. Lett. 107, 041801.
13. P. Adamson *et al.* (2011) [MINOS Collaboration], Phys. Rev.Lett. 107, 181802, arXiv: 1108.0015 [hep-ex].
14. G. L. Fogli, *et al.* (2011), Phys. Rev. D84, 053007, arXiv: 1106.6028 [hep-ph]
15. F. P. An *et al.*, Phys. Rev. Lett. 108, 171803 (2012), arXiv:1203.1669 [hep-ex]
16. J. K. Ahn *et al.*, *Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment*, arXiv:1204.0626 [hep-ex]
17. D. V. Forero, M. Trtola, J. W. F. Valle, Phys. Rev. D 86, 073012 (2012), arXiv:1205.4018 [hep-ph]
18. G. L. Fogli, *et al.* (2011), Phys. Rev. D 86, 013012 (2012), arXiv:1205.5254 [hep-ph]
19. M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado, and T. Schwetz, JHEP **12**, 123 (2012), arXiv:1209.3023 [hep-ph]
20. G. Aad *et al.* [The ATLAS Collaboration], Phys. Lett. B 716 (2012) 1; S. Chatrchyan *et al.* [The CMS Collaboration], Phys. Lett. B 716 (2012) 30.
21. A. Zee, *Phys. Lett. B* **93**, 389 (1980).
22. A. Zee, *Nucl. Phys. B* **264**, 99 (1986).
23. K. S. Babu, Phys. Lett. B **203**, 132 (1988).
24. S. Baek, P. Ko, and E. Senaha, *Can Zee-Babu model implemented with scalar dark matter explain both Fermi/LAT 130 GeV γ -ray excess and neutrino physics ?*, arXiv:1209.1685 [hep-ph] (2012).
25. K. S. Babu, C. Macesanu, Phys. Rev. D67 (2003) 073010, arXiv: 0212058 [hep-ph]
26. D. Aristizabal Sierra, M. Hirsch, JHEP 0612, 052 (2006), arXiv: 0609307 [hep-ph]
27. T. Araki and C. Q. Geng, Phys. Lett. B **694**, 113 (2010).
28. M. Nebot, J. F. Oliver, D. Palao, and A. Santamaria, Phys. Rev. D **77**, 093013 (2008).
29. D. Schmidt, T. Schwetz, H. Zhang, *Status of the Zee-Babu model for neutrino mass and possible tests at a like-sign linear collider*, arXiv:1402.2251 [hep-ph]
30. J. Herrero-Garcia, M. Nebot, N. Rius, A. Santamaria, *The Zee-Babu Model revisited in the light of new data*, arXiv:1402.4491 [hep-ph]

- 31. K. L. McDonald and B. H. J. McKellar (2003), hep-ph/0309270.
- 32. P. V. Dong and H. N. Long, *Surface Integral of Babu diagram*, [arXiv: hep-ph/0509007].
- 33. DAYA-BAY Collaboration, F. P. An et al, Phys. Rev. Lett. 108, 171803 (2012), arXiv:1203.1669
- 34. E. Ma, Phys. Rev. D **86**,117301 (2012)
- 35. S. F. King, Phys. Lett. B 659, 244 (2008), arXiv:0710.0530 [hep-ph].
- 36. D. Aristizabal Sierra, I. de Medeiros Varzielas, E. Houet, Phys. Rev. D 87.093009 (2013), arXiv:1302.6499 [hep-ph]